

21st-Century Precipitation Changes over the Los Angeles Region

Part III of the “Climate Change in the Los Angeles Region” Project



UCLA Institute of the Environment
and Sustainability

LARC
Los Angeles Regional Collaborative
for Climate Action and Sustainability

21st-Century Precipitation Changes over the Los Angeles Region

Part III of the “Climate Change in the Los Angeles Region” Project

Neil Berg
Graduate Student
UCLA Dept. of Atmospheric and Oceanic Sciences

Alex Hall
Professor
UCLA Dept. of Atmospheric and Oceanic Sciences

Fengpeng Sun
Assistant Researcher
UCLA Dept. of Atmospheric and Oceanic Sciences

Scott Capps
Assistant Researcher
UCLA Dept. of Atmospheric and Oceanic Sciences

Daniel Walton
Graduate Student
UCLA Dept. of Atmospheric and Oceanic Sciences

Baird Langenbrunner
Graduate Student
UCLA Dept. of Atmospheric and Oceanic Sciences

David Neelin
Professor
UCLA Dept. of Atmospheric and Oceanic Sciences

Support for this work was provided by the City of Los Angeles and the US Department of Energy as part of the American Recovery and Reinvestment Act of 2009. Additional funding was provided by the National Science Foundation (Grant #EF-1065863, "Collaborative Research: Do Microenvironments Govern Macroecology?") and the Southwest Climate Science Center. Additional support was provided by the National Science Foundation Grant AGS AGS-1102838. We thank two anonymous reviewers for their constructive comments.

For information, contact:

Professor Alex Hall
UCLA Dep't of Atmospheric and Oceanic Sciences
7955 Math Sciences Building
405 Hilgard Ave., Box 951565, Los Angeles, CA 90095
alexhall@atmos.ucla.edu • www.atmos.ucla.edu/csrl

Electronic copies of this study are available at: www.c-change.la

Table of Contents

Abstract.....	4
1. Introduction	5
2. Downscaling Techniques and Validation Results	8
2a. Dynamical Downscaling	8
1) Dynamical Downscaling Framework	8
2) Model Evaluation	11
2b. Hybrid Dynamical-Statistical Downscaling Framework.....	14
1) Empirical Orthogonal Function Analysis.....	14
2) Predicting Mode 1 Loadings	16
3) Validating Statistical Downscaling Techniques.....	17
4) Added Value Over Bilinear Interpolation.....	19
3. Dynamical-Statistical Downscaling Results	20
3a. Mid-21st-Century Changes	20
3b. End-of-21st-Century Changes	21
3c. Physical Mechanisms.....	22
4. Connection to Interannual Variability	24
4a. Context of Current Interannual Variability.....	24
4b. Relationship Between Future Climate Changes and Interannual Variability	25
5. Concluding Remarks	27
Figures	33
Tables	45
References	47

Abstract

A new hybrid dynamical-statistical downscaling technique is described to project mid- and end-of-21st-century local precipitation changes associated with 36 global climate models (GCMs) in Phase 5 of the Coupled Model Intercomparison Project archive over the greater Los Angeles region. Land-averaged precipitation changes, ensemble-mean changes, and the spread of those changes for both time slices are presented. It is demonstrated that the results are similar to what would be produced if expensive dynamical downscaling techniques were instead applied to all GCMs. Changes in land-averaged ensemble-mean precipitation are near zero for both time slices, reflecting the region's typical position in the models at the node of oppositely-signed large-scale precipitation changes. For both time slices, the intermodel spread of changes is only about 0.2–0.4 times as large as natural interannual variability in the baseline period. A caveat to these conclusions is that interannual variability in the tropical Pacific is generally regarded as a weakness of the GCMs. As a result, there is some chance the GCM responses in the tropical Pacific to a changing climate and associated impacts on Southern California precipitation are not credible. It is subjectively judged that this GCM weakness increases the uncertainty of regional precipitation change, perhaps by as much as 25%. Thus the possibility cannot be excluded that significant regional adaptation challenges related to either a precipitation increase or decrease would arise. However, the most likely downscaled outcome is a small change in local mean precipitation compared to natural variability, with large uncertainty in the sign of the change.

1. Introduction

Fresh water in the Los Angeles region comes from local storms, snowpack drainage, and groundwater. Identifying how climate change may impact these sources is of pressing concern for ecosystems and municipal, agricultural, and recreational purposes. In this study we aim to quantify 21st century climate change impacts to mean local sources of precipitation across the greater Los Angeles Region. Local sources contribute approximately 10% to the water supply in the city of Los Angeles (Villaraigosa 2008). However, in some areas, such as the San Fernando Valley, it contributes a larger portion (Sheng and Wilson 2008, ULARA 2011). Furthermore, these local sources may come under increasing pressure in the future (Erb et al. 2011). We do not address potential changes to imported water sources (e.g. the Colorado River) or extreme events (e.g. Das et. al 2013) in this study. A separate study will examine responses of local snowpack to climate change.

Projecting future precipitation changes over the Los Angeles region is challenging for two reasons. First, in GCM projections the region typically lies at the boundary of two oppositely-signed, large-scale zones of predicted precipitation change (van Oldenborgh et al. 2014), as described by the “rich-get-richer” or “wet regions get wetter and dry regions drier” effect (Chou and Neelin 2004, Held and Soden 2006, Trenberth 2011, Durack et al. 2012). Northern, midlatitude areas are projected to get wetter, while southern, sub-tropical areas are projected to become drier. Second, the complex topography of Southern California creates variations in precipitation that cannot be represented by coarse resolution GCM simulations. It is particularly important to adequately represent the coastal mountains over Southern California as they generally lead to significant orographic precipitation effects (Hughes et al. 2008, Neiman et al. 2002).

To address the limitations of coarse resolution GCMs, a common practice is to downscale global projections to a much finer resolution. Dynamical and statistical downscaling techniques are available to perform such a task. Dynamical downscaling solves the equations of motion and other atmospheric equations numerically, using a regional model that is forced along the boundaries by GCM output. This may represent the most physically consistent method to downscale climate data, although systematic biases may still be present in the downscaled simulation. The major tradeoff for dynamical downscaling is the expense of huge computational costs. Dynamical downscaling of climate change signals has been done for Southern California. For example, Duffy et al. (2005) dynamically downscaled two GCM projections, finding no statistically significant change in precipitation over Southern California.

Statistical downscaling is computationally cheap compared with dynamical downscaling, but hinges on currently existing relationships that may or may not hold true in the future. This technique has also been applied in the region of interest. For example, Hayhoe et al. (2004) statistically downscale four GCMs using historically derived empirical relationships and find small decreases in future wintertime precipitation in Southern California for three of the four simulations. A recent study by Pierce et al. (2012) uses separate dynamical and statistical downscaling techniques across 16 global climate models to examine future precipitation changes over California. Like Hayhoe et al. (2004), the statistical downscaling approaches used in Pierce et al. (2012) rely only on historical relationships (i.e. they assume stationarity) between variables when calculating climate change signals. After averaging across all downscaled projections, the authors find wintertime precipitation decreases of 5% over Southern California. Maurer (2007) statistically downscale future global precipitation and temperature output to drive a hydrologic model and find slight increases in wintertime precipitation over a basin in Southern California.

Das et al. (2013) statistically downscale 16 GCMs over the Sierra Nevada and find increased 3-day flood discharges, even though models tended to disagree on the sign of mean annual precipitation change. Pierce et al. (2013) also examine possible changes to daily precipitation over California. Using both dynamical and statistical downscaling techniques, they find evidence of increased wintertime precipitation over California, particularly over the northern part, due to an increase in daily precipitation intensity. Note that these previous studies rely on CMIP3 models, while this study only analyzes CMIP5 models. The two ensembles may exhibit different behavior in some cases. For example, Neelin et al. (2013) find that ensemble-mean drying in the CMIP3 archive is stronger over Southern California than in the CMIP5 archive.

The present study uses a new blended dynamical-statistical approach to project mid- and end-of-21st century December-January-March-February (DJFM) precipitation changes at a high resolution over the Los Angeles region. Whereas previous studies use only a dynamical or empirical statistical downscaling technique, this study develops statistical relationships directly from dynamically downscaled output. Using this method we are able to limit the assumption of stationarity that is often employed in statistical downscaling exercises (e.g. Hayhoe et al. 2004, Maurer 2007, Pierce et al. 2012). This technique also allows for downscaling of 36 GCMs in the CMIP5 archive (Table 1), providing analyses on intermodel spread and ensemble-mean changes. In addition to projecting 21st century precipitation changes over Southern California, another major aim of this study is to place climate change signals in context of the region's significant hydroclimate variability. Huge interannual variability in precipitation over Southern California is largely attributed to its relationships with large-scale natural climate variability patterns such as the El Niño–Southern Oscillation and the Pacific/North American Pattern (Cayan and Roads

1984, Redmond and Koch, 1991, Dettinger et al. 1998, Cayan et al. 1999, Leung et al. 2003, Berg et al. 2013).

The structure of the study is as follows: Section 2 describes the downscaling techniques and provides observational evaluation of the current climate simulation. Section 3 shows future precipitation changes according to 36 downscaled GCMs and explains the physical mechanisms behind the changes. A discussion of the relationship between climate change and interannual variability patterns is presented in Section 4, with a summary of major findings in Section 5.

2. Downscaling techniques and validation results

a. Dynamical downscaling

1) Dynamical downscaling framework

A dynamical downscaling simulation over Southern California was performed using the Weather Research and Forecasting Model (WRF), version 3.2 (WRF, Skamarock et al. 2008). We use three nested domains (18 km, 6 km, and 2 km) to reach a resolution high enough to represent the complex topography and coastlines of Southern California adequately. The three domains and topography associated with the outermost (18 km) domain are presented in Figure 1a. The outermost domain encompasses all of California and the adjacent Pacific Ocean, while the middle domain focuses on Southern California, including the southern Sierra Nevada mountain range. Finally, the innermost (2 km) domain is centered over the greater Los Angeles region. Topography associated with this domain is seen in Figure 1b.

Model sensitivity experiments were performed to find an optimal WRF configuration. Specifically, this simulation uses the Kain-Fritsch (new Eta) cumulus scheme (Kain 2004), Yonsei University boundary layer scheme (Hong et al. 2006), Purdue Lin microphysics scheme

(Lin et al. 1983), Rapid Radiative Transfer Model longwave radiation (Mlawer et al. 1997), Dudhia shortwave radiation schemes (Dudhia 1989), and the Noah land surface model (Chen and Dudhia 2001). Each domain has 43 sigma-levels in the vertical and vertical resolution is increased below 3 km to better simulate surface and boundary layer processes.

Two time periods are simulated to initially project mid-21st century precipitation changes. We focus first on a “baseline” period spanning 1981–2000. In this case, WRF is forced along the boundaries of the outermost domain by the North American Regional Reanalysis (NARR). Then we simulate a range of future climates based on model output from five CMIP5 GCMs (CCSM4, CNRM-CM5, GFDL-CM3, MIROC-ESM-CHEM, and MPI-ESM-LR), all under the RCP8.5 emissions scenario. For each future simulation, baseline (1981–2000) boundary conditions from NARR are perturbed with future monthly climatological changes (2041–2060 average minus 1981–2000 average) to atmospheric variables and imposed on WRF. 3-dimensional atmospheric variables that were perturbed include temperature, relative humidity, zonal and meridional winds, and geopotential heights. Surface temperature, relative humidity, winds, and pressure were also perturbed. This technique has been used previously (e.g., Schär et al. 1996, Hara et al. 2008, Knutson et al. 2008, Kawase et al. 2009, Lauer et al. 2010, Rasmussen et al. 2011, Seo and Xie 2011 and Gutmann et al. 2012) and estimates future climates as perturbations to the same baseline mean-state, corresponding roughly to the present day. A limitation to this technique is that future interannual variability equals that of the baseline period. Implications of this limitation when analyzing downscaled changes are discussed further in sections 3c and 5. For an application of this downscaling method applied to future warming over the Los Angeles region, the reader is referred to Sun et al. (2014).

We first perform a 20-year future simulation (2041–2060), downscaling climate change signals in CCSM4. Computational expenses prevent full 20-year simulations for other models, so we perform a sensitivity test examining how long of a future period we needed to simulate to capture the full 20-year climate change signal. Figure 2 shows that by only simulating three future years (2058–2060) we are able to capture the full 20-year signal to a high degree of accuracy. (Other consecutive 3-year periods between 2041–2060 may also be highly representative of the full 20-year time span, though computational resources prevented this analysis). Spatial structures between the two signals are tightly correlated, with only slight discrepancies seen in the coastal zone. Averaged over the land, the 20-year and 3-year signals are -46.7 and -46.6 mm/wet season, respectively. Relying on this knowledge, we next dynamically downscale the four other GCMs (CNRM-CM5, GFDL-CM3, MIROC-ESM-CHEM, and MPI-ESM-LR) for 2058–2060. In each simulation, boundary conditions were created by adding the 2041–2060 minus 1981–2000 GCM changes to the 1998–2000 NARR values. Therefore, interannual variability over 2058–2060 is the same as 1998–2000; however, the perturbations imposed in the future runs represent a climate change signal associated with much longer averaging periods. Statistical downscaling techniques are then developed based on these 2058–2060 minus 1998–2000 dynamically downscaled changes (section 2.b). The reader is directed to the supplementary material for a discussion on possible biases in the statistical model due to interannual variability differences between the 3-year-long and 20-year-long changes. Namely, it is found that the dynamical model underestimates the magnitude of average precipitation changes by around 20% due to being based on 3-year-long signals (2058–2060 minus 1998–2000) compared to the 20-year-long signals (2041–2060 minus 1981–2000). Thus statistical

estimates based on these changes may be associated with a reduced spread by a complementary amount. The implications of this error are examined further in section 5.

2) Model evaluation: spatial and temporal variability in the baseline

Before presenting the results of the climate change experiments, we compare simulated interannual precipitation variations in the baseline (1981–2000) 2-km WRF output to observations. We use three observational datasets: California Irrigation Management Information System (CIMIS, <http://wwwcimis.water.ca.gov/>), NOAA Climate Prediction Center 0.25°x0.25° Daily US UNIFIED Precipitation (CPC, <http://www.esrl.noaa.gov/psd/data/gridded/data.unified.html>), and the 0.5°x0.5° gridded University of Delaware Precipitation product (UDel, http://www.esrl.noaa.gov/psd/data/gridded/data.UDel_AirT_Precip.html). Correlations between these data sets and WRF output may be less than 1.0 for multiple reasons, including WRF inaccuracies, unresolved sub-grid scale topography (i.e., elevation mismatch between the location being sampled and the WRF grid cell average), poor observational data quality, and inaccuracies in the boundary conditions (NARR) forcing WRF. Assuming the observational products are perfect, the model evaluation serves as a test of WRF's ability to reproduce precipitation variations over the Los Angeles region when coarse resolution conditions (NARR) are imposed on it. If WRF is able to transform this coarse-resolution data into regional climate information that closely matches accurate observational products, we are confident WRF can regionalize the GCM signal in a way that is consistent with the real atmosphere's dynamics.

In Fig. 3a, we correlate monthly DJFM precipitation accumulations in the baseline period between each CIMIS station and the nearest WRF grid point. Each correlation is based on a

maximum sample size of 80 (4 wet-season months X 20 baseline years = 80 values). However, there are missing values in the observations, leading to an average sample size of 45 values. Twelve of the thirteen stations have correlations to WRF above 0.5, and more than half have correlations above 0.7. Thus, WRF generally simulates monthly precipitation variations at rain gauges across the domain reasonably well. The lone exception is Santa Barbara ($r=0.37$). We speculate that WRF simulates the complex interactions between small-scale circulations and rainfall at this location of intense coastal topography poorly. In Fig. 3b, we correlate 1981–2000 DJFM-mean precipitation accumulations (20 values per grid point) between each CPC grid point and the nearest corresponding WRF grid point. Correlations greater than 0.6 are found across nearly the entire domain, with very high values ($r>0.9$) found along much of the densely populated coastal region. The domain-average correlation is 0.82. Thus interannual variability simulated in WRF and that recorded in the CPC gridded product is very similar.

Additional validation of precipitation variability in the baseline WRF simulation is presented in Figure 4. This figure compares interannual variability of monthly precipitation amounts in the three observational datasets (CIMIS, CPC, and UDel) and WRF output at the scale of the domain. Each white, gray, or black dot in Fig. 4 represents monthly precipitation accumulations for each of the 20 baseline years that are simulated. The large dots represent monthly climatologies for each dataset. Two comparisons can be made in Fig. 4. The first is between CIMIS station-averaged monthly precipitation accumulations (white dots, see Fig. 3a for station locations) and corresponding accumulations averaged over the nearest grid points in the 2 km WRF domain (light gray dots). The levels of interannual variability in CIMIS and WRF station-averages are very similar for each month, and the two time series are highly correlated ($r=0.88$). Climatological accumulations for each month are also very similar, with an average

monthly climatology difference between the two datasets of approximately 6 mm, or 8%.

Particularly noteworthy is the similarity between the observed and modeled bimodal structure of the temporal precipitation distribution, seen most dramatically in January and February. Both datasets capture the extremely dry (<25 mm) and wet (>250 mm) months within the baseline period.

The second comparison to make in Fig. 4 is between the UDel, CPC, and WRF land-average monthly accumulations (medium gray, dark gray, and black dots, respectively). Like the CIMIS comparison, WRF variability in monthly precipitation accumulations tightly matches what is observed in the UDel (average $r=0.94$) and CPC (average $r=0.96$) datasets. Differences in monthly climatologies between WRF and UDel are approximately 17 mm (28%), and approximately 9 mm (15%) between WRF and CPC. Interestingly, for both WRF-based and observation-based datasets, there are strong similarities in magnitude between the station-averaged (white and light gray dots) and land-averaged values (medium gray, dark gray, and black dots). This indicates that the station-averages adequately sample the land fraction of the domain. For example, the average monthly climatology difference between CIMIS station-averaged (white dots) and CPC land-averaged (dark gray dots) values is only approximately 16 mm.

Finally, we assess WRF's ability to simulate spatial variations in station-averaged (in the case of CIMIS rain gauges) and land-averaged (in the case of UDel and CPC gridded observations) precipitation totals over the baseline period. Results are seen in Figure 5, which shows scatter plots between simulated and observed (CIMIS: black circles, UDel: red circles, CPC: cyan circles) station or land-averaged wet-season total accumulations. Note that CIMIS observations begin in 1989, so only 12 wet seasons are included in this portion of the plot. WRF

reproduces the CIMIS observations ($r=0.83$, average bias of +15 mm) better than UDel ($r=0.59$, bias of +229 mm) or CPC ($r=0.55$, bias of +221 mm). The large disagreement between WRF and the two gridded products is likely due to the horizontal resolution differences between them. Coarse resolutions in the gridded products ($0.25^\circ \times 0.25^\circ$ for CPC and $0.5^\circ \times 0.5^\circ$ for UDel) may not resolve the full orographic effects on precipitation, which are included in WRF and of course the station measurements. As noted above, discrepancies between WRF and CIMIS values or any data product may arise due to sub-grid scale topography and poor observational data quality, in addition to model deficiencies and inaccuracies in the boundary conditions (NARR).

b. Hybrid dynamical-statistical downscaling framework

1) Empirical orthogonal function analysis

Here we present the hybrid dynamical-statistical approach to generating future precipitation projections. We begin by forming statistical relationships between precipitation changes in the five dynamically downscaled GCMs to large-scale parameters in GCM output. The first step is identifying common spatial patterns between monthly wet-season precipitation changes (2058–2060 minus 1998–2000) for all five downscaled models. Each GCM's dynamically downscaled monthly precipitation changes over the course of the wet season (DJFM) can be seen in Figure 6. We make two remarks on the variations in Fig. 6. First, there is variation in the sign and magnitude of mid-21st-century precipitation changes in dynamically downscaled results. Some downscaled GCMs, such as CCSM4 (Fig. 6, first row), show future drying over most of the coastal zone and high elevations for all months, while others, such as CNRM-CM5 (Fig. 6, second row), project moistening for much of the domain over most months. Other outcomes lie between these two cases, and are not necessarily consistent in sign across the domain. Second,

we note that although there is large variation across downscaled models and months, there appears to be a common area where *most* of the action occurs—a pattern tied to orography, with enhanced loading in the coastal zone and throughout the mountainous regions. This suggests that performing an empirical orthogonal function (EOF) analysis on the aggregated set of these monthly precipitation change patterns could yield a single, robust spatial pattern of change.

Following this reasoning, an EOF analysis is performed over the spatial patterns in Fig. 6. Since the EOF analysis spans both models and months, the patterns it generates maximize both intermodel and intermonthly variability. The three leading modes are shown in Figure 7. The first accounts for 70% of the variability seen in Fig. 6, confirming our suspicion that the majority of the variance can be accounted for with a single spatial pattern. Mode 1 physically represents the dominant orographic pattern of precipitation over Southern California (Hughes et al. 2008, Conil and Hall 2005, Neiman et al. 2002). Other precipitation patterns, such as “blocked events” (Hughes et al. 2008, Neiman et al. 2002), may be somewhat represented in modes 2 and 3. A corresponding 20-value (5 dynamically downscaled models x 4 months) series of mode 1 loadings is also produced from the EOF analysis. These loadings represent the contribution of the spatial pattern of mode 1 to each model’s monthly precipitation change. Since this mode accounts for the majority of intermodel and intermonthly variability, it should be possible to “predict” the dynamically downscaled precipitation changes in Fig. 6 with reasonable accuracy simply by multiplying the spatial pattern of mode 1 by each model’s monthly mode 1 loading. (While modes 2 and 3 may represent a physical phenomenon associated with precipitation change, we ignore them due to the small variance that is captured in each mode, 7% and 5%, respectively.) Blending the statistical methods of an EOF analysis and dynamical downscaled simulations forms what we call a hybrid dynamical–statistical downscaling technique. For an

example of how this blended statistical–dynamical downscaling approach can be applied to regional warming patterns, the reader is referred to Walton et al. (2014).

2) Predicting mode 1 loadings

We have calculated mode 1 loadings for the five dynamically downscaled models, but we need a method for predicting the mode 1 loadings for the other GCMs if they were dynamically downscaled. The first step is to relate the known mode 1 loadings to a large-scale predictor variable available from the GCMs, in this case precipitation. In Figure 8a, we correlate mid-21st-century monthly DJFM precipitation changes over the north Pacific in the five GCMs that were dynamically downscaled to the loading series associated with mode 1. Each GCM is regridded to a common horizontal resolution ($1.5^\circ \times 1.5^\circ$) before performing the correlation. A dipole correlation pattern emerges. GCM precipitation change over the Gulf of Alaska shows anticorrelations to regional precipitation changes associated with mode 1, while the Pacific Ocean adjacent to California shows positive correlations. A physical interpretation of this correlation pattern is discussed in section 4b. We tried several statistical techniques to relate mode 1 loadings to GCM precipitation changes, including single and multivariable linear regression and a projection-based dot-product technique. The strongest (highest correlations) and most robust (across the downscaled GCMs) relationship was found using linear regression, where mode 1 loadings are predicted by two predictor variables: GCM precipitation changes averaged over the two regions spanning the dipole correlation pattern (black boxes, Fig. 8a). This yields a single equation to predict a given GCM's mode 1 loading, if that GCM were dynamically downscaled, based only on its mid-21st-century precipitation change across the northeast Pacific Ocean. A caveat is that these predictive equations hinge on the training set of

dynamically downscaled models, in this case CCSM4, CNRM-CM5, GFDL-CM3, MIROC-ESM-CHEM, and MPI-ESM-LR. A different set of models could give different relationships between GCM and local precipitation changes. However, some robustness is provided to the predictive relationships by developing them on a set of GCMs that span the dry-to-wet parameter space (Figs. 6 and 10).

3) Validating statistical downscaling techniques

The statistical model may capture dynamical model output imperfectly for two reasons: (1) mode 1 is an imperfect representation of regional precipitation change, and (2) it is impossible to predict mode 1 loadings perfectly. Knowing the loadings associated with mode 1 from our EOF analysis of dynamically downscaled simulations, we can test how accurate DJFM-mean changes are based solely on mode 1, i.e. the first source of error. This comparison is shown in Figure 9. Recall that the EOF analysis is performed over monthly changes, so DJFM-mean values shown here are calculated by averaging individual monthly patterns to produce a seasonal mean. First we compare the spatial patterns between the dynamically downscaled changes (Fig. 9a) and those based on mode 1 (Fig. 9b). In general the spatial patterns are very well correlated, aside from modest discrepancies in the Mojave Desert regions. WRF (y-axis) versus mode 1-based (x-axis) precipitation changes from Figs. 9a and 9b, now averaged over land, are scattered in Fig. 9c. Mode 1 captures the land-averaged precipitation change extremely well, with the mode 1 changes and the WRF changes falling almost perfectly on the line $y=x$. These results confirm that if we have perfect knowledge of mode 1 loadings, then statistically downscaled ensemble-mean changes and the spread in these changes are highly representative of the corresponding

dynamically downscaled changes. This is especially true when considering the change averaged over the region's land areas.

Next we analyze the errors associated with imperfect predictions of mode 1 loadings, i.e., the second source of error in the statistical model, using cross-validation experiments. These experiments use differing subsets of the five dynamically downscaled output to develop a predictive equation for mode 1 loadings. We then predict mode 1 loadings for all dynamically downscaled models and compare them to the actual loadings. Specifically, we perform five experiments. The experiment number is equal to the number of dynamically downscaled models used to determine mode 1 loadings. Each experiment is performed for a varying number of trial runs, consistent with the number of ways it is possible to combine the models. For example, experiment 1 uses one model set of DJFM monthly precipitation changes to determine mode 1 loadings (i.e. any one row in Fig. 6). It has five trials since there are five possible DJFM monthly change values that can be used to predict mode 1 loadings. Experiment 2 uses two model sets of DJFM monthly changes to predict mode 1 loadings for all models, yielding 10 unique combinations (i.e. any two rows in Fig. 6). Experiments 3 (i.e. any three rows in Fig. 6) and 4 (i.e. any four rows in Fig. 6) have 10 and five trials, respectively, and experiment 5 (all rows in Fig. 6), has only one trial.

In essence, we are testing the robustness of the statistical model as more and more dynamically downscaled information is included in its training. For each trial run in each experiment, we perform all analyses described in section 2b.2 for the dynamically downscaled models being used for mode 1 predictions. That is, we first perform an EOF analysis over the spatial patterns of monthly precipitation changes (e.g. 4 patterns per trial in experiment 1). The EOF analysis yields a series of mode 1 loadings, which are then correlated to the corresponding

GCM mid-21st-century precipitation changes across the Pacific Ocean. Finally, GCM mid-21st-century precipitation changes over the regions of maximum positive and negative correlation (which varies according to each trial's correlation map, but is similar to Fig. 8a for all trials) are regressed against that trial's mode 1 loadings. This yields a predictive equation for mode 1 loadings for each of the five dynamically downscaled models, which can be compared to the known mode 1 loadings.

Table 2 summarizes the uncertainty of the statistical model due to errors in the predictions of mode 1 loadings. The error averaged over all models for all trials is shown in the right column. Errors decrease steadily as the number of models used in the EOF analysis increases. This makes sense, since more intermonthly, intermodel variability is included as more information is fed into the analyses. Specifically, average error is reduced from over 100% when using just one or two models, to just 13% when using five models. It can appear that using four models gives a smaller percent error (-2%) than when five models are used (-13%). However, this simply reflects an average over a very large range when four models are used (-483% to +466%) compared to a much smaller range using five models (-103% to +137%).

4) Value added over bilinear interpolation

Here we justify the development of our hybrid dynamical-statistical downscaling technique by comparing results to a simple bilinear regression of the raw GCM data down to 2 km. Figure 10 provides evidence that the hybrid downscaling technique adds significant value in spatial patterns compared to bilinearly interpolating GCM data over Southern California. For each GCM in Fig. 10, spatial patterns that emerge in the interpolated results are broad in scale and have no way of capturing the leading spatial pattern seen in the dynamical downscaling associated with

orographic effects. Land-averaged changes between the interpolated and dynamically downscaled GCMs can be quite similar, and in some cases closer (likely fortuitous given the coarse GCM resolution) than the corresponding statistically downscaled changes. However, orographic influences on precipitation (e.g. Hughes et al. 2008) are simply not captured in either the raw or interpolated GCM data. Conversely, the hybrid dynamical-statistical downscaling technique is able to capture the orographic imprint on precipitation changes with reasonable accuracy. It should also be noted that the standard deviation between the statistically and dynamically downscaled land-averaged changes is 6.5 and 9.5 mm/wet season, respectively. Thus the statistical model may underestimate the spread of changes on the order of 30%. We will assess the implications of this potential error in Section 4a.

3. Dynamical-statistical downscaling results

Here we predict the regional precipitation projections for all 36 GCMs (Table 1), using the statistical model described in the previous section.

a. Mid-21st-century changes

Mid-21st-century DJFM-mean precipitation changes from all 36 downscaled GCMs are shown in Figure 11. Recall that the downscaled projections in Fig. 11 are forced to have the same spatial pattern (that of mode 1, Fig. 7) and that the spatial pattern is dialed up or down based on the predicted loading for that GCM. Precipitation changes projected using full dynamical downscaling would have somewhat more spatial heterogeneity than those shown in Fig. 11. Thus we do not focus on the spatial patterns of change, but rather interpret results from a land-average

perspective. The land-average can be predicted by the statistical model with a high degree of accuracy once mode 1 loadings are known (see Section 2b).

Fig. 11 shows an apparently large range of projected changes across models. 13 models project increased precipitation (average of +8.1 mm/wet season) and 23 models project decreased precipitation (average of -8.5 mm/wet season). The most extreme models are MIROC5 and IPSL-CM5A-MR, which project changes of approximately +19 and -25 mm/wet season across the land, respectively. The ensemble-mean land-average change is -2.5 mm/wet season, reflecting a large degree of cancellation between moistening and drying tendencies. Note that the statistical model may underestimate the spread of changes up to 50% (section 2), so the true intermodel variability of changes may be 50% larger than described here.

b. End-of-21st-century changes

The statistical model can also be used to project end-of-century (2081-2100 – 1981-2000) precipitation changes. As seen by the dark blue dots in Figure 4, the ensemble-mean change is near zero for each month and the spread of those changes is smaller than current levels of variability, similar to the mid-century case. In addition to downscaled changes, we also present interpolated GCM changes in Fig. 4 (light blue dots). Like the mid-21st-century changes, the ensemble-mean change by the end of the 21st century is near zero for each month. Taken as a whole, Fig. 4 indicates that the model-average downscaled and interpolated GCM scenario for the Los Angeles region is very little precipitation change throughout the 21st century.

c. Physical mechanisms

Recall that the dynamically downscaled future simulations are based on GCM perturbations to NARR climatologies along the lateral boundaries to 3-dimensional temperature, moisture, winds, and geopotential heights, and surface winds, moisture, and pressure. Interpreting the physical mechanisms behind precipitation changes are constrained to this methodology. While storms entering the outermost domain in WRF (Fig. 1a) are structurally identical between the baseline and future simulations (a limitation raised in Rasmussen et al. 2011), the perturbation method allows for storms to evolve differently as they propagate towards the innermost domain. For example, possible changes to large-scale circulations, such as the jet stream, can be captured in the 3-dimensional wind and geopotential height perturbations. Moisture content in future storms could also change due to perturbations in relative humidity. Future storm strength could also be modified by perturbations to surface pressure and 3-dimensional geopotential heights. Bearing this information in mind, we revisit Figure 8 to identify the physical mechanisms underpinning downscaled precipitation changes (Fig. 11).

As described in section 2b.2, Fig. 8a shows that precipitation changes over Los Angeles are related to large-scale precipitation changes over extreme northern and north/central portions of the eastern Pacific Ocean. The patterns in Fig. 8a suggest that average jet stream position changes across the Pacific Ocean are largely controlling precipitation changes over Los Angeles. A recent study by Neelin et al. (2013) analyzed the relationship between end-of-century California December-January-February (DJF) precipitation changes and 200 mb zonal wind speed changes over the northeast Pacific Ocean in 15 CMIP5 GCMs [cf. Fig. 1, Neelin et al. (2013)]. Precipitation changes over the California land-ocean region are found to be significantly related to changes in the jet stream (i.e. 200 mb zonal winds) and associated storm tracks. GCMs

projecting increased jet stream wind speeds, associated with an eastward and poleward jet extension, tend to steer more storms toward the coast and lead to overall precipitation increases in this region. GCMs that show weak eastward jet extension and/or wind speed enhancement are associated with minimal precipitation changes. Specifically, the authors find a correlation of 0.76 between end-of-century DJF precipitation changes over California and 200 mb zonal wind speed over a certain region of the northwest Pacific.

Though our domain of interest is the Los Angeles region rather than the whole state of California, we follow the arguments presented in Neelin et al. (2013), and perform an analysis relating GCM 200 mb zonal wind speed changes to downscaled precipitation changes. 200 mb zonal wind speed changes (2041–2060 minus 1981–2000) for the 36 downscaled models are correlated at each grid point in the GCM domain to the domain-averaged downscaled precipitation changes. Each GCM is regridded to a common horizontal resolution ($1.5^\circ \times 1.5^\circ$) before performing the correlation. The results are shown in Fig. 8b. Strong negative correlations are seen across most of the Gulf of Alaska and into western Canada. Conversely, strong positive correlations are seen across the entire north central Pacific Ocean, centered on Hawaii. This dipole pattern echoes the results found in Neelin et al. (2013) and indicates how jet stream positioning and strength influence future precipitation over the Los Angeles region. Specifically, GCMs that project regional increases/decreases in jet stream strength off the coast of Southern California lead to increased/decreased precipitation over Los Angeles.

4. Connection to interannual variability

a. Context of current interannual variability

Here we place the intermodel spread of future precipitation changes in the context of the region's natural precipitation variability. Examining Fig. 4, we compare the variability across statistically downscaled model projections of future changes (red dots) and levels of interannual variability for the wet-season (black dots). Averaged across each month, the standard deviations for the downscaled mid-century precipitation changes are 15, 15, 12, and 14 mm/wet season, respectively. (The standard deviations of end-of-century values are very similar.) The standard deviation of baseline interannual variability of WRF land-averaged, monthly-averaged accumulations (black dots, Fig. 4) is 61 mm/wet season. Thus, the intermodel variations of downscaled future changes in average precipitation are roughly 25% of the current interannual variability. As noted in Section 2, the statistical model may underestimate the standard deviation of the precipitation changes, due to imperfect knowledge of mode 1 loadings, probably by about 30%. So potentially the true standard deviation of precipitation changes is roughly 40% of the variability. But even after factoring in this possible bias, it is clear that the interannual precipitation variability is large compared to potential changes in the mean. Of course, the mean changes would be sustained on time scales much longer than a year, potentially leading to adaptation challenges. For example, the downscaled models with the most extreme drying and moistening tendencies are associated with mean precipitation changes on the order of 10%. However, such challenges would only materialize if the more extreme models are correct; the average downscaled and interpolated GCM outcome is virtually no precipitation change for the entire century.

b. Relationship between future climate changes and interannual variability

So far we have argued that GCM placement of jet stream and storm tracks in the north Pacific Ocean is the main driver of intermodel variability in future precipitation changes over Los Angeles. Previous studies have also shown jet stream placement, strength, and storm track steering over the Pacific Ocean can shift due to natural climate variability patterns (Chen and van den Dool 1997, Straus and Shukla 1997, Held et al. 1989). These jet stream and storm track shifts impact the amount of precipitation over Southern California (Berg et al. 2013, Athanasiadis et al. 2010). The importance of the jet stream for future precipitation change suggests a tight link between the physical underpinnings of interannual variability and simulated climate change.

We begin addressing the relationship between interannual and intermodel variability by analyzing baseline DJFM precipitation from the 1981–2000 WRF simulation forced by NARR. An EOF analysis is performed over 20 spatial patterns of DJFM-averaged precipitation anomalies corresponding to each year of the baseline simulation. The patterns are calculated as anomalies relative to the 1981–2000 DJFM climatology. The leading mode accounts for 86% of the variability, and the corresponding spatial pattern is very similar to the first mode of intermodel variability determined from the climate change experiments (Fig. 12). The leading modes of variability in both the baseline and future cases reflect the strong orographic enhancement of precipitation and the influence of blocking in the coastal zone across the greater Los Angeles region (Hughes et al. 2008). After performing the EOF analysis over the baseline precipitation fields, we then correlate the time series associated with mode 1 (Fig. 12a) to 1981–2000 precipitation anomalies at each grid point in the NARR data. These correlation coefficients are plotted in Figure 8c, and can be compared to the future case (Fig. 8a, section 3). Both cases

show a tongue of positive correlations that extend from the coast of California westward into the Pacific Ocean. This tongue is then flanked on the north and south by large swaths of anticorrelations. We also perform a correlation between baseline precipitation and 200 mb zonal wind anomalies in the NARR data (Fig. 8d) and compare it to the corresponding case associated with future changes in the GCMs (Fig. 8b, section 3c). Both cases show a dipole pattern of large positive correlations across the southern half of the eastern Pacific Ocean and large negative correlations in the northern half.

Such similarities in Figure 8 confirm that the dynamics of baseline interannual variability are nearly identical to those underpinning future intermodel uncertainty. That is, the region's precipitation currently vacillates between wet and dry periods with a pattern heavily modulated by orography. The vacillations are largely due to natural variations in the position and strength of the jet stream and subsequent storm track steering. Models that tend to deflect the jet stream and storms away from Southern California yield drier climates in the future, while models showing a tendency toward jet stream strengthening and increased storm activity over Southern California project a wetter climate. Thus the collection of moistening and drying tendencies in the CMIP5 ensemble can likely be understood as an "excitation" of a natural mode of variability.

5. Concluding remarks

This study uses a hybrid dynamical-statistical downscaling technique to examine mid- and end-of-21st-century precipitation changes over the greater Los Angeles region under the RCP8.5 emissions scenario. Modeling dynamically downscaled precipitation changes with statistical methods, we downscale 36 GCMs in the CMIP5 archive based on changes in each model's large-scale precipitation fields. There are three major findings of this study. First, the ensemble-mean change for both time slices is essentially zero. Second, while downscaled CMIP5 models disagree on both the sign and magnitude of future precipitation changes over Los Angeles, the spread of possible changes is modest compared to current levels of variability. For both time slices, the statistical model estimates that the standard deviation of land-averaged precipitation change is about 0.2 to 0.25 of the standard deviation of the interannual variability. As shown in section 2, the statistical model may underestimate the intermodel spread by as much as 30% due to imperfect knowledge of mode 1 loadings. So the true standard deviation of the precipitation change, if all GCMs were downscaled dynamically, could be closer to 0.4 of the interannual variability standard deviation. Thus even after allowing for potential error in the statistical model, current shifts between wet and dry years are greater than average changes in even the most extreme model projections. However, the sustained moistening or drying seen in the most extreme statistically downscaled models could lead to adaptation challenges. Though these changes are unlikely, they amount to roughly 10% changes in mean precipitation for both time slices. Finally, robust similarities are found between the intermodel variability of future changes and interannual variability of baseline precipitation anomalies. Jet stream placement and strength currently dictates winter precipitation amounts, and also dictates the sign and magnitude of future precipitation changes. To the degree there is uncertainty in future precipitation change

over the Los Angeles region, it is due to differences in the simulated response of this phenomenon to anthropogenic forcing.

While there is a great opportunity to assign probabilities of future changes based on an ensemble of projection outcomes, no single method perfectly accomplishes this task. Ensemble-mean forecasts have proven skillful in producing most likely outcomes for climate variations on the seasonal time scale (e.g. Doblus-Reyes et al. 2003, Palmer et al. 2005b), along with hurricane paths and other weather patterns (e.g. Zhang and Krishnamurti 1998, Krishnamurti et al. 2000). Interpreting an ensemble-mean projection as the most likely outcome for 21st-century precipitation, however, is complicated due in part to the fact that the range of outcomes in the ensemble spans positive and negative changes (van Oldenborgh et al. 2014, their Fig. A1.6). Our result of near-zero ensemble-mean change simply reflects offsetting tendencies of moistening and drying in the GCMs. It could be misleading not to acknowledge that some change, either positive or negative, is likely to occur. At the same time, for nearly all projections, the magnitude of the change is small compared to natural variability. In this sense we interpret the *most likely* scenario as a small change in precipitation compared to natural variability, with large uncertainty on the sign of the change.

A critique to this probabilistic estimate is that we are weighting each projection equally in the ensemble, assuming the quality of each model is the same (Tebaldi and Knutti 2007, Knutti 2010, Shepherd 2014). One way to address this issue is to weight each model based on its ability to simulate historical precipitation or the large-scale circulations controlling precipitation over California. Swain et al. (2014) identified 12 CMIP5 GCMs that best simulate historical 500 mb geopotential height over the northeastern Pacific Ocean according to reanalysis distributions. Geopotential height in this region strongly influences California's precipitation (Fig. 8). By

examining downscaled projections from these 12 “good” models in Fig. 11, we again find that precipitation changes are modest compared to natural variability, and that the sign of the change is uncertain. Specifically, 8 models project increased precipitation, 4 project decreased precipitation, and the ensemble-mean change is +2.2 mm/wet season. (Similar results between the 12- and 36-model ensembles is perhaps not too surprising as Langford et al. (2014) found that most CMIP5 models have reasonable skill in simulating California precipitation.) This simple analysis indicates that the major conclusions of this study are not sensitive to which models are included in the ensemble.

Our result of near-zero ensemble-mean precipitation change over Los Angeles can be interpreted in terms of the well-accepted understanding of global precipitation change whereby patterns of precipitation become enhanced, such that wet regions become wetter and dry regions become drier (Chou and Neelin 2004, Neelin et al. 2006, Held and Soden 2006). This leads to increased precipitation over convection zones and drying outside of the convection zones. On average, Southern California is positioned between areas dominated by these competing tendencies: increased precipitation to its north in the mid-latitudes and decreased precipitation to the south within the subtropics. However, in some GCMs the region is north of the boundary between the two zones, while in others it is south of it. As such, precipitation projections over this region tend to negate one another and yield small ensemble-mean projections.

One interesting finding from this study is that intermodel variability between the statistically downscaled (red dots, Fig. 4) changes is approximately half the size of the variability according to the GCM-interpolated changes (pink dots). We attribute this spread suppression in the statistical model to two sources. First, inaccurate mode 1 predictions by the statistical model (section 2) can underestimate the intermodel spread by at least 30%. Second, statistical

relationships are derived from 3-year-long dynamically downscaled changes (2058–2060 minus 1998–2000). On average, 1998–2000 was found to have somewhat lower precipitation than 1981–2000. This has the effect of reducing the magnitude of statistically-estimated climate change signals, and the spread associated with them, likely about 20% (Figure S1, Table S1). If these two error sources were eliminated, the spread would probably increase by about 50%—therefore approximately accounting for the difference between statistically downscaled and GCM-interpolated inter-model variability seen in Fig. 4. We also note that the results in this study are limited to our choice of WRF as the regional model. Unlike larger regional modeling efforts such as CORDEX (e.g., Nikulin et al. 2012) and NARCCAP (e.g., Wang et al. 2009), computational costs prevented an examination of how sensitive this study’s results are to multiple regional model simulations. However, we found strong agreement between our results and those from an independent study that statistically downscaled precipitation changes over the greater Los Angeles region (personal communication with Lee Alexanderson, 2014, Los Angeles County Department of Public Works).

Differences between the regional model outcomes and those of the GCMs may also stem from our method of perturbing baseline boundary conditions using future climatological changes. For example, one could instead directly downscale raw historical and future GCM data to calculate changes, as opposed to perturbing baseline conditions derived from reanalysis. We are currently conducting research to test whether this direct method gives different results from downscaling changes in the climatology through a perturbation to reanalysis-based boundary conditions.

Given the agreement between the GCMs and the downscaled information in the ensemble-mean outcome, it seems unlikely that a different dynamical downscaling technique would generate a systemically different answer. The hybrid dynamical-statistical downscaling technique could be applied beyond the Los Angeles region. It may be especially appropriate in areas that share these two characteristics with the domain of interest in our study: (1) changes in the large-scale circulation govern precipitation change, allowing for development of credible GCM scaling factors, and (2) local precipitation changes are heavily influenced by orography, leading to diagnosed local response patterns, as encapsulated by the leading EOF patterns. Thus it would be applicable for any mid-to-high latitude location with significant topography.

An important caveat relating to the El Niño–Southern Oscillation (ENSO) phenomenon applies to the conclusions of this study. In the current climate, ENSO influences the position of the Northern Hemisphere jet stream and storm tracks across the eastern Pacific Ocean through atmospheric teleconnections (Held et al. 1989, Chen and van den Dool 1997, Straus and Shukla 1997). These shifts have a statistically-detectable effect on precipitation over Southern California. During La Niña events, the jet tends to move northward towards the Gulf of Alaska, leading to drier than average conditions across Southern California. Under El Niño conditions, the jet tends to extend south and eastward, steering storms more directly across southern regions of US, including Southern California (Redmond and Koch 1991, Dettinger et al. 1998, Cayan et al. 1999, Leung et al. 2003, Berg et al. 2013). The CMIP5 ensemble of GCMs has shown improvements in the simulation of ENSO compared to the CMIP3 ensemble, particularly in the amplitude and time scale of the phenomenon. However, the CMIP5 models still exhibit significant errors, especially in the irregularity of the phenomenon and its spatial pattern (Flato et al. 2013). A detailed examination of the implications of these tropical Pacific errors for

precipitation change over Southern California is beyond the scope of this study. However, it seems possible that the GCM projections of future ENSO behavior may be affected by them.

If these errors were corrected, modestly different outcomes for Southern California precipitation might result, owing to the link between ENSO variability and Southern California precipitation. When an ENSO event occurs, it accounts for roughly 2/3 of the variance in Southern California precipitation. However, only about 40% of wet seasons can be considered strong ENSO events (Schonher and Nicholson, 1989). Thus roughly one quarter of the variance of Southern California precipitation can be traced to ENSO. The remaining three-quarters of the variance is linked to shifts of the jet stream unrelated to tropical Pacific variability, similar to those portrayed in Fig. 8d, and which are also the mechanism generating intermodal spread in the CMIP5 ensemble. While ENSO is a mechanism generating regional precipitation variability, it is not the most important. ENSO errors in the GCMs may introduce somewhat more uncertainty in our regional precipitation projections than what is implied by the downscaled intermodel spread alone. It is impossible to quantify this effect precisely with present knowledge, but the role ENSO currently plays in Southern California precipitation does at least offer a useful guide. We estimate that ENSO GCM errors increase the uncertainty by an amount approximately proportional to the fraction of the variance ENSO accounts for in current climate—by about 25%. This additional uncertainty underscores the need for regional planning that allows for a variety of future precipitation change outcomes.

Figures




FIG. 1. (a) 18 km – 6 km – 2 km WRF domains and 18 km topography; (b) 2 km domain and topography. Black lines in (a) and (b) show US state boundaries and Los Angeles County for reference. Also seen in (b) are the Channel Islands. Topography is color contoured every 200 m in both (a) and (b).




FIG. 2. Dynamically downscaled CCSM4 climate change signals according (a) 20-year (2041-2060 – 1981-2000) and (b) 3-year (2058-2060 – 1998-2000) changes. Difference between the 20-year and 3-year signals is shown in (c). Topography is contoured every 750 m in thin black lines. Unit is mm/wet season.




FIG. 3. (a) Correlation coefficients of monthly 1981-2000 DJFM accumulated precipitation between CIMIS stations and the nearest grid point in the 2 km WRF output. Topography is contoured every 750 m in thin black lines. (b) Correlation coefficients between 1981-2000 DJFM-mean accumulated precipitation amounts between CPC grid cells and nearest corresponding WRF grid cells.




FIG. 4. Monthly precipitation accumulations (mm/month) averaged over CIMIS stations (white dots), WRF-NARR grid points nearest to CIMIS stations (light great dots), land-averaged in the UDel observational dataset (medium grey dots), land-averaged in the CPC observational dataset (dark grey dots), and land-averaged in the WRF-NARR output (black dots). Larger dots in each case represent monthly climatologies. Also shown are monthly mid- and end-of-21st century precipitation *changes* (mm/wet season) relative to the base-period climate according to 36 statistically downscaled (red/blue) and interpolated (pink/light blue) CMIP5 GCMs. Larger red/blue and pink/light blue dots represent ensemble-mean monthly changes.




FIG 5. Scatter plots between simulated and observed wet-season (DJFM) climatological precipitation over the baseline period (1981-2000). Black circles show CIMIS station-averaged amounts vs. averages over the nearest WRF-NARR grid points, red circles show land-averaged UDel vs. WRF-NARR values, and cyan circles show land-averaged CPC vs. WRF values. The line $y=x$ is shown as a solid black line. Unit is mm/wet season.




FIG. 6. DJFM monthly precipitation changes (2058-2060 minus 1998-2000) for each dynamically downscaled GCM. Blue shading indicates moistening, red shading indicates drying. Unit is mm/wet season. Topography is contoured every 750 m in thin black lines.




FIG 7. Leading three modes of variability based on EOF analysis of spatial patterns seen in Fig. 6. Mode 1 accounts for 70% of the variability, mode 2 accounts for 7%, and mode 3 accounts for 5%.




FIG. 8. Correlation coefficients between (a) mid-21st century monthly DJFM precipitation changes (2041-2060 minus 1981-2000) according to the five dynamically downscaled GCMs and the time series associated with EOF 1 (Fig. 7). Black squares represent averaging area of GCM precipitation to predict EOF 1 loadings. (b) domain-averaged downscaled precipitation changes (Fig. 11) and corresponding mid-21st century 200 mb zonal wind speed changes for all available models, (c) monthly DJFM precipitation anomalies in the 1981-2000 NARR data and time series associated with EOF 1 over that time period (Fig. 12a), and (d) domain-averaged 1981-2000 precipitation anomalies and corresponding 200 mb zonal wind speed anomalies in the NARR data.




FIG. 9. (a) Dynamically-downscaled DJFM-mean precipitation changes for each model. (b) Mode 1-based DJFM-mean precipitation changes for each model. (c) Scatter plot comparing domain-averaged DJFM-mean changes from WRF (y-axis) and mode-1 (x-axis), with the line $x=y$ shown as a solid black line. Unit in each plot is mm/wet season.




FIG. 10. Comparison of mid-21st century precipitation changes for the downscaled models according to the respective raw GCM data (first column), bilinearly interpolated GCM data to 2 km (second column), the hybrid statistical-dynamical downscaling technique (third column), and the dynamical downscaling (fourth column). Land-averaged changes (mm/wet season) are reported in the top right of each panel.




FIG. 11. Downscaled mid-21st century precipitation changes according to 36 GCMs. Blue shading indicates future moistening, while brown shading indicates future drying. Unit is mm/wet season.




FIG. 12. Leading modes of precipitation variability over the baseline (a) and future (b), same as Fig. 7a). Baseline precipitation anomalies are calculated relative to the 1981-2000 climatology. Future changes are calculated as 2058-2060 – 1998-2000. See text for details.

Tables

Model	Institute
ACCESS1-0	Commonwealth Scientific and Industrial Research Organization
ACCESS1-3	Commonwealth Scientific and Industrial Research Organization
BNU-ESM	College of Global Change and Earth System Science, Beijing Normal University
CCSM4	National Center for Atmospheric Research
CESM1-BGC	National Science Foundation, Department of Energy, National Center for Atmospheric Research
CESM1-CAM5	National Science Foundation, Department of Energy, National Center for Atmospheric Research
CMCC-CESM	Euro-Mediterranean Center of Climate Change
CMCC-CM	Euro-Mediterranean Center of Climate Change
CMCC-CMS	Euro-Mediterranean Center of Climate Change
CNRM-CM5	Centre National de Recherches Meteorologiques
CSIRO-Mk3-6-0	Commonwealth Scientific and Industrial Research Organization
CanESM2	Canadian Centre for Climate Modeling and Analysis
EC-EARTH	EC-Earth Consortium
FGOALS-g2	LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences
GFDL-CM3	NOAA Geophysical Fluid Dynamics Laboratory
GFDL-ESM2G	NOAA Geophysical Fluid Dynamics Laboratory
GFDL-ESM2M	NOAA Geophysical Fluid Dynamics Laboratory
GISS-E2-H	NASA Goddard Institute for Space Studies
GISS-E2-R	NASA Goddard Institute for Space Studies
HadGEM2-AO	Met Office Hadley Centre
HadGEM2-CC	Met Office Hadley Centre
HadGEM2-ES	Met Office Hadley Centre
IPSL-CM5A-LR	Institut Pierre Simon Laplace
IPSL-CM5A-MR	Institut Pierre Simon Laplace
IPSL-CM5B-LR	Institut Pierre Simon Laplace
MIROC-ESM	AORI (U. Tokyo), NIES, JAMSTEC
MIROC-ESM-CHEM	AORI (U. Tokyo), NIES, JAMSTEC
MIROC5	AORI (U. Tokyo), NIES, JAMSTEC
MPI-ESM-LR	Max Planck Institute for Meteorology
MPI-ESM-MR	Max Planck Institute for Meteorology
MRI-CGCM3	Meteorological Research Institute
NorESM1-M	Norwegian Climate Center
NorESM1-ME	Norwegian Climate Center
bcc-csm1-1	Beijing Climate Center, China Meteorological Administration
bcc-csm1-1-m	Beijing Climate Center, China Meteorological Administration
inmcm4	Institute for Numerical Mathematics

TABLE 1. List of CMIP5 models and corresponding institutions used in this study.

Number of models (number of trials)	Average/min/max percent error between actual and predicted mode 1 loadings (%)
1 (5)	-160 / -1831 / 1981
2 (10)	-139 / -1366 / 473
3 (10)	-72 / -1048 / 558
4 (5)	-2 / -483 / 466
5 (1)	-13 / -103 / 137

TABLE 2. Quantifying the error associated with imperfect predictions of mode 1 loadings in the statistical model using a cross-validation exercise. Number of models used and the number of unique combinations (“trials”) of those models (i.e. any row in Fig. 6) are presented in the left column. The average, maximum and minimum percent error averaged over all models for all trials and is seen in the right column.

References

- Athanasiadis, P.J. and J.M. Wallace, 2010: Patterns of Wintertime Jet Stream Variability and Their Relation to the Storm Tracks. *J. Atmos. Sci.*, **67**, 1361–1381.
- Berg, N., A. Hall, S.B. Capps, and M. Hughes, 2013: El Nino–Southern Oscillation Impacts on Winter Winds over Southern California. *Clim. Dyn.*, DOI 10.1007/s00382-012-1461-6
- Cayan, D.R. and J.O. Roads, 1984: Local Relationships between United States West Coast Precipitation and Monthly Mean Circulation Parameters. *Mon. Wea. Rev.*, **112**, 1276–1282.
- Cayan, D.R., K.T. Redmond, and L.G. Riddle, 1999: ENSO and hydrologic extremes in the Western United States. *J. Climate*, **12**, 2881–2893.
- Chen, F., and J. Dudhia, 2001: Coupling an advanced land surface-hydrology model with the Penn State-NCAR MM5 modeling system. Part I: Model implementation and sensitivity. *Mon. Wea. Rev.*, **129**, 569-585.
- Chen, W.Y., and H.M. van den Dool, 1997: Asymmetric impact of tropical SST anomalies on atmospheric internal variability over the North Pacific. *J. Atmos. Sci.*, **54**, 725–740.
- Chou, W.Y. and J.D. Neelin, 2004: Mechanisms of global warming impacts on regional tropical precipitation. *J. Climate*, **17**, 2688-2005.
- Conil S., and A. Hall, 2005: Local Regimes of Atmospheric Variability: A Case Study of Southern California. *J. Climate*, **19**, 4308-4325.
- Das T., E.P. Maurer, D.W. Pierce, M.D. Dettinger, and D.R. Cayan, 2013: Increases in flood magnitudes in California under warming climates. *J. Hydrology*, **501**, 101-110.
- Dettinger, M.D., D.R. Cayan, H.F. Diaz, and D.M. Meko, 1998: North–south precipitation patterns in Western North America on interannual-to-decadal timescales. *J. Climate*, **11**, 3095–3111.
- Doblas-Reyes, F. J., V. Pavan, and D. B. Stephenson, 2003: The skill of multimodel seasonal forecasts of the wintertime North Atlantic Oscillation. *Clim. Dynamics.*, **21**, 501-504.
- Dudhia, J., 1989: Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model. *J. Atmos. Sci.*, **46**, 3077-3107.
- Duffy, P.B., R.W. Arritt, J. Coquard, W. Gutowski, J. Han, J. Iorio, J. Kim, L.-R. Leung, J. Roads, and Z. Zeledon, 2005: Simulations of Present and Future Climates in the Western United States with Four Nested Regional Climate Models. *J. Climate*, **19**, 873–895.
- Durack, P.J., S.E. Wijffels, and R.J. Matear, 2012: Ocean salinities reveal strong global water cycle intensification during 1950 to 2000. *Science*, **336**, 455-458.

Erb, T., Tam W., Kharaghani S., Hanna M., Poresky A., and P. Reidy, 2011: Stormwater as a Resource: Rainwater Harvesting in the City of Los Angeles. Proceedings of the Water Environment Federation, WEFTEC 2011: Session 101 through Session 110 , pp. 6935–6947(13)

Flato, G., J. Marotzke, B. Abiodun, P. Braconnot, S.C. Chou, W. Collins, P. Cox, F. Driouech, S. Emori, V. Eyring, C. Forest, P. Gleckler, E. Guilyardi, C. Jakob, V. Kattsov, C. Reason and M. Rummukainen, 2013: Evaluation of Climate Models. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley (eds)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Gutmann, E., R. Rasmussen, C. Liu, D.J. Gochis, and M. Clark, 2012: A comparison of statistical and dynamical downscaling of winter precipitation over complex terrain. *J. Climate*, **25**, 262–281.

Hara, M., T. Yoshikane, H. Kawase, and F. Kimura, 2008: Estimation of the impact of global warming on snow depth in Japan by the pseudo-global-warming method. *Hydrol. Res. Lett.*, **2**, 61–64

Hayhoe K., D. Cayan, C.B. Field, P.C. Frumhoff, E.P. Maurer, N.L. Miller, S.C. Moser, S.H. Schneider, K.N. Cahill, E.E. Cleland, L. Dale, R. Drapeki, R.M. Hanemann, L.S. Kalkstein, J. Lenihan, C.K. Lunch, R.P. Neilson, S.C. Sheridan, and J.H. Verville, 2004: Emissions pathways, climate change, and impacts on California, *PNAS*, **34**, 12422–12427.

Held, I.M., S.W. Lyons, and S. Nigam, 1989: Transients and the extratropical response to El Nino. *J. Atmos. Sci.*, **46**, 163–174.

Held, I.M. and B.J. Soden, 2006: Robust Responses of the Hydrological Cycle to Global Warming. *J. Climate*, **19**, 5686–5699.

Hong, S.-Y., Y. Noh, and J. Dudhia, 2006: A new vertical diffusion package with an explicit treatment of entrainment processes. *Mon. Wea. Rev.*, **134**, 2318-2341.

Hughes, M., A. Hall, and R.G. Fovell, 2008: Blocking in Areas of Complex Topography, and Its Influence on Rainfall Distribution. *J. Atmos. Sci.* **66**, 508–518.

Kain, J. S., 2004: The Kain-Fritsch convective parameterization: An update. *J. Appl. Meteor. Soc.*, **43**, 170-181.

Kawase, H., T. Yoshikane, M. Hara, F. Kimura, T. Yasunari, B. Ailikun, H. Ueda, and T. Inoue, 2009: Intermodel variability of future changes in the Baiu rainband estimated by the pseudo global warming downscaling method. *J. Geophys. Res.*, **114**, D24110. doi:10.1029/2009JD011803.

- Knutsen, T. R., J. J. Sirutius, S. T. Garner, G. A. Vecchi, and I. M. Held, 2008: Simulated reduction in Atlantic hurricane frequency under twenty-first century warming conditions. *Nat. Geosci.*, **1**, 359–364.
- Knutti, R., 2010: The end of model democracy? *Climatic Change*, **102**, 395–404.
- Krishnamurti, T. N., C. M. Kishtawal, Z. Zhang, T. LaRow, D. Bachiochi, E. Williford, S. Gadgil, and S. Surendran, 2000: Multimodel Ensemble Forecasts for Weather and Seasonal Climate. *J. Climate*, **13**, 4196–4216.
- Langford, S., S. Stevenson, and D. Noone, 2014: Analysis of Low-Frequency Precipitation Variability in CMIP5 Historical Simulations for Southwestern North America. *J. Climate*, **27**, 2735–2756.
- Lauer, A., K. Hamilton, Y. Wang, V. T. J. Phillips, and R. Bennartz, 2010: The impact of global warming on marine boundary layer clouds over the eastern Pacific—A regional model study. *J. Climate*, **23**, 5844–5863.
- Leung, L.R., Y. Qian, X. Bian, and A. Hunt, 2003: Hydroclimate of the Western United States Based on Observations and Regional Climate Simulations of 1981–2000. Part II: Mesoscale ENSO Anomalies. *J. Climate*, **16**, 1912–1928.
- Lin, Y.-L., R. D. Farley, and H. D. Orville, 1983: Bulk parameterization of the snow field in a cloud model. *J. Climate Appl. Meteor.*, **22**, 1065–1092.
- Maurer, E. P. 2007: Uncertainty in hydrologic impact of climate change in the Sierra Nevada, California, under two emissions scenarios. *Climatic Change*, **82**, 309–325.
- Mlawer, E. J., S. J. Taubman, P. D. Brown, M. J. Iacono, and S. A. Clough, 1997: Radiative transfer for inhomogeneous atmosphere: RRTM, a validated correlated-k model for the longwave. *Journal of Geophysical Research: Atmospheres* (1984–2012), **102** (D14), 16663–16682.
- Neelin, J.D., M. Munnich, H. Su, J.E. Meyerson, and C.E. Holloway, 2006: *PNAS*. **16**, 6110–6115.
- Neelin, J.D., B. Langenbrunner, J.E. Meyerson, A. Hall, and N. Berg, 2013: California Winter Precipitation Change under Global Warming in the Coupled Model Intercomparison Project Phase 5 Ensemble. *J. Climate*. **00**, 1–19.
- Neiman, P.J., F.M. Ralph, A.B. White, D.E. Kingsmill, and P.O.G. Persson, 2002: The Statistical Relationship between Upslope Flow and Rainfall in California’s Coastal Mountains: Observations during CALJET. *Mon. Wea. Rev.*, **130**, 1468–1492.
- Nikulin, G., C. Jones, F. Giorgi, G. Asrar, M. Büchner, R. Cerezo-Mota, O. B. Christensen, M. Déqué, J. Fernandez, A. Hänsler, E. van Meijgaard, P. Samuelsson, M. Bamba Sylla, and L. Sushama, 2012: Precipitation Climatology in an Ensemble of CORDEX-Africa Regional

Climate Simulations. *J. Climate*, **25**, 6057–6078. doi: <http://dx.doi.org/10.1175/JCLI-D-11-00375.1>

Palmer, T. N., F. J. Doblas-Reyes, R. Hagedorn, and A. Weisheimer, 2005b: Probabilistic prediction of climate using multi-model ensembles: from basics to applications. *Phil. Trans. R. Society.*, B 360: 1991-1998.

Pierce, D. W., D. R. Cayan, T. Das, E. P. Maurer, N. L. Miller, Y. Bao, M. Kanamitsu, K. Yoshimura, M. A. Snyder, L. C. Sloan, G. Franco, and M. Tyree, 2013: The Key Role of Heavy Precipitation Events in Climate Model Disagreements of Future Annual Precipitation Changes in California. *J. Climate*, **26**, 5879-5896.

Pierce, D.W., T. Das, D.R. Cayan, E.P. Maurer, N.L. Miller, Y. Bao, M. Kanamitsu, K. Yoshimura, M.A. Snyder, L.C. Sloan, G. Franco, and M. Tyree, 2012: Probabilistic estimates of future changes in California temperature and precipitation using statistical and dynamical downscaling. *Clim. Dyn.*, DOI:10.1007/s00382-012-1337-9

Rasmussen, R. and Coauthors, 2011: High-Resolution Coupled Climate Runoff Simulations of Seasonal Snowfall over Colorado: A Process Study of Current and Warmer Climate. *J. Climate*, **24**, 3015–3048

Redmond, K.T. and R.W. Koch, 1991: Surface climate and streamflow variability in the Western United States and their relationship to large-scale circulation indices. *Water Resour. Res.*, **27**, 2381–2399.

Schär, C., C. Frei, D. Lüthi, and H. C. Davies, 1996: Surrogate climate-change scenarios for regional climate models. *Geophys. Res. Lett.*, **23**(6), 669–672. doi:10.1029/96GL00265.

Schonher, T. and S.E. Nicholson, 1989: The Relationship between California Rainfall and ENSO Events. *J. Climate*, **2**, 1258-1269.

Seo, H., and S.-P. Xie, 2011: Response and impact of equatorial ocean dynamics and tropical instability waves in the tropical Atlantic under global warming: A regional coupled downscaling study. *J. Geophys. Res.*, **116**, C03026. doi:10.1029/2010JC006670.

Sheng J, and J.P. Wilson, 2008: An Implimentation Blueprint for Enhancing Groundwater Recharge in the San Fernando Basin, California: University of Southern California GIS Research Laboratory Technical Report No. 7

Shepherd T. G., 2014: Atmospheric circulation as a source of uncertainty in climate change projections. *Nature Geoscience*, **7**, 703-708.

Skamarock, W.C., Klemp J.B., Dudhia J., Gill D.O., Barker D.M., Duda M.G., Huang X-Y, Wang W., Powers J.G., 2008: A Description of the Advanced Research WRF Version 3. NCAR Technical Note, NCAR/TN-475+STR.

Straus, D.M., and J. Shukla, 1997: Variations of midlatitude transient dynamics associated with ENSO. *J Atmos. Sci.*, **54**, 777–790.

Sun, F., D. Walton, and A. Hall, 2014: A hybrid dynamical–statistical downscaling technique, part II: End-of-century warming projections predict a new climate state in the Los Angeles region. Submitted to *J. Climate*.

Swain, D. L., M. Tsiang, M. Haugen, D. Singh, A. Charland, B. Rajaratnam, and N. Diffenbaugh, 2014: The Extraordinary California Drought of 2013/2014: Character, Context, and the Role of Climate Change [in "Explaining Extremes of 2013 from a Climate Perspective"]. *Bull. Amer. Meteor. Soc.*, **95** (9), S3-S7.

Tebaldi C. and R. Knutti, 2007: The use of the multi-model ensemble in probabilistic climate projections. *Phil. Trans R. Soc.*, **365**, 2053-2075.

Trenberth, K.E. 2011: Changes in precipitation with climate change. *Climate Res.*, **47**, 123-138.

ULARA Watermaster, 2013: Annual Report of the Upper Los Angeles River Area Watermaster for the 2011-2012 water year.

van Oldenborgh, G.J., M. Collins, J. Arblaster, J.H. Christensen, J. Marotzke, S.B. Power, M. Rummukainen and T. Zhou 2014: Atlas of global and regional climate projections. *Climate Change 2013: The Physical Sciences Basis*. [Stocker, T.F. et al. (eds.)]. Cambridge University Press, 1311-1394.

Villaraigosa, A.R., 2008: Securing L.A.'s Water Supply, City of Los Angeles Water Supply Action Plain, City of Los Angeles Department of Water and Power.
http://www.greencitiescalifornia.org/assets/water/LA_Emergency-Water-Conservation-Plan_Water-Supply-Report-2008.pdf

Walton D., Sun F., A. Hall, and S. Capps, 2014: A Hybrid Dynamical-Statistical Downscaling Technique, Part I: Development and Validation of the Technique. *Submitted to J. Climate*.

Wang, S.-Y., R. R. Gillies, E. S. Takle, and W. J. Gutowski Jr. 2009: Evaluation of precipitation in the Intermountain Region as simulated by the NARCCAP regional climate models, *Geophys. Res. Lett.*, 36, L11704, doi:[10.1029/2009GL037930](https://doi.org/10.1029/2009GL037930).

Zhang, Z. and T. N. Krishnamurti 1998: A Perturbation Method for Hurricane Ensemble Prediction. *Mon. Wea. Rev.*, **127**, 447-469.